


$$\textcircled{1} \iint_M (x^2 + y^2 - y) dx dy, \quad M: x^2 + y^2 \leq 4, x \leq 0$$




$$x = r \cdot \cos \varphi$$

$$y = r \cdot \sin \varphi$$

$$x^2 + y^2 = r^2$$


$$\frac{\pi}{2} \leq \varphi \leq \frac{3\pi}{2}$$

$$0 \leq r \leq 2$$




$$\begin{aligned} & \int_{\frac{\pi}{2}}^{\frac{3\pi}{2}} \left[\int_0^2 (r^2 - r \cdot \sin \varphi) \cdot r dr \right] d\varphi \\ &= \int_{\frac{\pi}{2}}^{\frac{3\pi}{2}} \left[\int_0^2 (r^3 - r^2 \cdot \sin \varphi) dr \right] d\varphi \\ &= \int_{\frac{\pi}{2}}^{\frac{3\pi}{2}} \left[\frac{r^4}{4} - \frac{r^3}{3} \sin \varphi \right]_0^2 d\varphi = \int_{\frac{\pi}{2}}^{\frac{3\pi}{2}} \left(4 - \frac{8}{3} \sin \varphi \right) d\varphi \\ &= \left[4\varphi + \frac{8}{3} \cos \varphi \right]_{\frac{\pi}{2}}^{\frac{3\pi}{2}} \\ &= 6\pi + \frac{8}{3} \cdot \underbrace{\cos \frac{3\pi}{2}}_{=0} - 2\pi - \frac{8}{3} \cdot \underbrace{\cos \frac{\pi}{2}}_{=0} = 4\pi \end{aligned}$$

② $\iint_R x \sqrt{x^2+y^2} dx dy$ $x^2+y^2 \geq 1$, $x^2+y^2 \leq 4$, $x \geq 0$, $y \geq 0$



$$0 \leq \theta \leq \frac{\pi}{2}$$

$$1 \leq r \leq 2$$



$$x = r \cdot \cos \theta$$

$$x^2 + y^2 = r^2$$

$$\int_0^{\pi/2} \left[\int_1^2 \underbrace{r \cdot \cos \theta \cdot \sqrt{r^2}}_{r^3 \cdot \cos \theta} \cdot r dr \right] d\theta$$

$$= \int_0^{\pi/2} \cos \theta d\theta \cdot \int_1^2 r^3 dr$$

$$= \left[\sin \theta \right]_0^{\pi/2} \cdot \left[\frac{r^4}{4} \right]_1^2$$

$$= \left(\sin \frac{\pi}{2} - \sin 0 \right) \cdot \left(4 - \frac{1}{4} \right) = \frac{15}{4}$$

$$=$$