



Second order linear differential equation Nonhomogeneous equation and Wronskian Interactive tests

Robert Mařík

July 14, 2006

ROBERT MAŘÍK
LDE and Wronskian
file ldr24.tex

Theory

Tests

Home Page

Print

Title Page



Page 1 of 15

Go Back

Full Screen

Close

Quit



Look at three or four or twenty my quizzes and
then fill in my _____ please!

1. Theory



ROBERT MAŘÍK
LDE and Wronskian
file ldr24.tex

Theory
Tests

Home Page

Print

Title Page

◀ ▶

◀ ▶

Page 2 of 15

Go Back

Full Screen

Close

Quit

Definition 1 (second order linear differential equation) Let p, q be real numbers, f be functions continuous on the interval I . The equation

$$y'' + py' + qy = f(x) \quad (1)$$

is said to be a **second order linear differential equation with constant coefficients**.

Definition 2 (special types of 2nd order LDE) Equation (1) is said to be **homogeneous** if $f(x) = 0$ for all $x \in I$ and **nonhomogeneous** otherwise.

Definition 3 (associated homogeneous equation) Consider nonhomogeneous equation (1). Homogeneous equation

$$y'' + py' + qy = 0. \quad (2)$$

with the left-hand side identical with equation (1) is called a **homogeneous equation associated to the nonhomogeneous equation (1)**.

Theorem 1 (general solution of nonhomogeneous second order LDE) Let $y_1(x)$ and $y_2(x)$ be fundamental system of solutions of the homogeneous LDE (2) and $y_p(x)$ be an arbitrary particular solution of the nonhomogeneous LDE (1). Then the function

$$y(x, C_1, C_2) = C_1 y_1(x) + C_2 y_2(x) + y_p(x), \quad C_1 \in \mathbb{R}, C_2 \in \mathbb{R} \quad (3)$$

is a general solution of the nonhomogeneous LDE (1).



Theorem 2 (solution of the nonhomogeneous LDE with constant coefficients)

Consider the second order LDE with constant coefficients

$$y'' + py' + qy = f(x). \quad (4)$$

Let $y_1(x)$ and $y_2(x)$ be a fundamental system of solutions of the associated homogeneous equation. Let $A(x)$ and $B(x)$ be differentiable functions with derivatives given by

$$A'(x) = \frac{W_1(x)}{W(x)} \quad \text{and} \quad B'(x) = \frac{W_2(x)}{W(x)},$$

where

$$W(x) = \begin{vmatrix} y_1(x) & y_2(x) \\ y_1'(x) & y_2'(x) \end{vmatrix}, \quad W_1(x) = \begin{vmatrix} 0 & y_2(x) \\ f(x) & y_2'(x) \end{vmatrix}, \quad W_2(x) = \begin{vmatrix} y_1(x) & 0 \\ y_1'(x) & f(x) \end{vmatrix}.$$

The function $y_p(x)$ defined by the relation

$$y_p(x) = A(x)y_1(x) + B(x)y_2(x) \quad (5)$$

is a particular solution of (4). The function

$$y(x) = C_1y_1(x) + C_2y_2(x) + y_p(x), \quad C_1 \in \mathbb{R}, C_2 \in \mathbb{R},$$

is a general solution of (4).



2. Tests

On the following pages you find some tests which guide you through the process of solving second order LDE via variation of constants.

- The functions from the fundamental system are not unique. If you choose the system as on the lectures, the correct answer is unique up to the order of the functions. This quiz can handle this answer.
- If your answer on the fundamental system is correct (or if you scratch your answer and asks for correct answer by pressing "?" button), follow the hint which tells you which one of the two functions is $y_1(x)$ and which one is $y_2(x)$. This hint is in the popup window as well as written in the test. Then the answers to the questions on W , W_1 , W_2 , A' and B' are unique.
- The correct answers on functions A and B are unique up to an additive constant.
- The particular solution is unique up to an additive factor which is a linear combination of the functions from the fundamental system. This test is able to grade the users answer on particular solution correctly. Thus both answers $y = 1$ and $y = 1 + \sin(x) + 2 \cos(x)$ are correct for the equation $y'' + y = 1$.
- In the field with general solution all functions $y = 1 + A \sin(x) + B \cos(x)$, $y = 1 + \sin(x) + A \cos(x) + 3B \sin(x)$ or $y = 1 + A \sin(x) - B(\cos(x) - \sin(x))$ are equivalent, as excepted from the fact that there are many possibilities how to write the general solution.

Home Page

Print

Title Page

⏪ ⏩

⏪ ⏩

Page 4 of 15

Go Back

Full Screen

Close

Quit



Quiz Solve $y'' + y = 1$.

1. Char. eq. of the assoc. hom. eq. (in variable z):
2. Fundamental system:
3. Look for the particular solution in the form $y_p = A(x)y_1(x) + B(x)y_2(x)$

$$W(x) = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} =$$
$$W_1(x) = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} =$$
$$W_2(x) = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} =$$

4. $A'(x) = \frac{W_1}{W} =$
 $B'(x) = \frac{W_2}{W} =$

5. Integrating we get

$$A(x) =$$

$$B(x) =$$

6. Particular solution: $y_p =$

7. Find general solution, use constants A and B .

$$y =$$



Quiz Solve $y'' + y = x$.

1. Char. eq. of the assoc. hom. eq. (in variable z):
2. Fundamental system:
3. Look for the particular solution in the form $y_p = A(x)y_1(x) + B(x)y_2(x)$

$$W(x) = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} =$$
$$W_1(x) = \begin{vmatrix} y_1 & x \\ y_1' & 1 \end{vmatrix} =$$
$$W_2(x) = \begin{vmatrix} x & y_2 \\ 1 & y_2' \end{vmatrix} =$$

4. $A'(x) = \frac{W_1}{W} =$
 $B'(x) = \frac{W_2}{W} =$

5. Integrating we get

$$A(x) =$$

$$B(x) =$$

6. Particular solution: $y_p =$

7. Find general solution, use constants A and B .

$$y =$$



Quiz Solve $y'' - 3y' + 2y = e^x$.

1. Char. eq. of the assoc. hom. eq. (in variable z):
2. Fundamental system:
3. Look for the particular solution in the form $y_p = A(x)y_1(x) + B(x)y_2(x)$

$$\begin{aligned} W(x) &= \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = \\ W_1(x) &= \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = \\ W_2(x) &= \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = \end{aligned}$$

4.
$$A'(x) = \frac{W_1}{W} =$$
$$B'(x) = \frac{W_2}{W} =$$

5. Integrating we get

$$A(x) =$$

$$B(x) =$$

6. Particular solution: $y_p =$

7. Find general solution, use constants A and B .

$$y =$$



Quiz Solve $y'' + 4y = x$.

1. Char. eq. of the assoc. hom. eq. (in variable z):
2. Fundamental system:
3. Look for the particular solution in the form $y_p = A(x)y_1(x) + B(x)y_2(x)$

$$W(x) = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} =$$
$$W_1(x) = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} =$$
$$W_2(x) = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} =$$

4. $A'(x) = \frac{W_1}{W} =$
 $B'(x) = \frac{W_2}{W} =$

5. Integrating we get

$$A(x) =$$

$$B(x) =$$

6. Particular solution: $y_p =$

7. Find general solution, use constants A and B .

$$y =$$



Quiz Solve $y'' + y = x + 2$.

1. Char. eq. of the assoc. hom. eq. (in variable z):
2. Fundamental system:
3. Look for the particular solution in the form $y_p = A(x)y_1(x) + B(x)y_2(x)$

$$\begin{aligned} W(x) &= \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = \\ W_1(x) &= \begin{vmatrix} y_1 & y_2 \\ y_1' & 0 \end{vmatrix} = \\ W_2(x) &= \begin{vmatrix} 0 & y_2 \\ y_1' & y_2' \end{vmatrix} = \end{aligned}$$

4.
$$A'(x) = \frac{W_1}{W} =$$
$$B'(x) = \frac{W_2}{W} =$$

5. Integrating we get

$$A(x) =$$

$$B(x) =$$

6. Particular solution: $y_p =$

7. Find general solution, use constants A and B .

$$y =$$



Quiz Solve $y'' - 2y' + y = \frac{e^x}{x}$.

1. Char. eq. of the assoc. hom. eq. (in variable z):
2. Fundamental system:
3. Look for the particular solution in the form $y_p = A(x)y_1(x) + B(x)y_2(x)$

$$\begin{aligned} W(x) &= \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = \\ W_1(x) &= \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = \\ W_2(x) &= \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = \end{aligned}$$

4.
$$A'(x) = \frac{W_1}{W} =$$
$$B'(x) = \frac{W_2}{W} =$$

5. Integrating we get

$$A(x) =$$
$$B(x) =$$

6. Particular solution: $y_p =$

7. Find general solution, use constants A and B .

$$y =$$



Quiz Solve $y'' - 5y' + 6y = e^{2x}$.

1. Char. eq. of the assoc. hom. eq. (in variable z):
2. Fundamental system:
3. Look for the particular solution in the form $y_p = A(x)y_1(x) + B(x)y_2(x)$

$$\begin{aligned} W(x) &= \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = \\ W_1(x) &= \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = \\ W_2(x) &= \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = \end{aligned}$$

4.
$$A'(x) = \frac{W_1}{W} =$$
$$B'(x) = \frac{W_2}{W} =$$

5. Integrating we get

$$A(x) =$$

$$B(x) =$$

6. Particular solution: $y_p =$

7. Find general solution, use constants A and B .

$$y =$$



Home Page

Print

Title Page

◀ ▶

◀ ▶

Page 12 of 15

Go Back

Full Screen

Close

Quit

Quiz Solve $y'' + y = \frac{\cos x}{\sin x}$.

1. Char. eq. of the assoc. hom. eq. (in variable z):
2. Fundamental system:
3. Look for the particular solution in the form $y_p = A(x)y_1(x) + B(x)y_2(x)$

$$W(x) = \begin{vmatrix} \\ \end{vmatrix} =$$

$$W_1(x) = \begin{vmatrix} \\ \end{vmatrix} =$$

$$W_2(x) = \begin{vmatrix} \\ \end{vmatrix} =$$

4. $A'(x) = \frac{W_1}{W} =$
 $B'(x) = \frac{W_2}{W} =$

5. Integrating we get
 $A(x) =$
 $B(x) =$

6. Particular solution: $y_p =$

7. Find general solution, use constants A and B .
 $y =$



Quiz Solve $y'' + 2y' - 3y = e^x$.

1. Char. eq. of the assoc. hom. eq. (in variable z):
2. Fundamental system:
3. Look for the particular solution in the form $y_p = A(x)y_1(x) + B(x)y_2(x)$

$$\begin{aligned} W(x) &= \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = \\ W_1(x) &= \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = \\ W_2(x) &= \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = \end{aligned}$$

4.
$$A'(x) = \frac{W_1}{W} =$$
$$B'(x) = \frac{W_2}{W} =$$

5. Integrating we get

$$A(x) =$$

$$B(x) =$$

6. Particular solution: $y_p =$

7. Find general solution, use constants A and B .

$$y =$$



Quiz Solve $y'' - 4y = e^{2x}$.

1. Char. eq. of the assoc. hom. eq. (in variable z):
2. Fundamental system:
3. Look for the particular solution in the form $y_p = A(x)y_1(x) + B(x)y_2(x)$

$$W(x) = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} =$$
$$W_1(x) = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} =$$
$$W_2(x) = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} =$$

4. $A'(x) = \frac{W_1}{W} =$
 $B'(x) = \frac{W_2}{W} =$

5. Integrating we get

$$A(x) =$$

$$B(x) =$$

6. Particular solution: $y_p =$

7. Find general solution, use constants A and B .

$$y =$$



Quiz Solve $y'' - 4y = xe^x$.

1. Char. eq. of the assoc. hom. eq. (in variable z):
2. Fundamental system:
3. Look for the particular solution in the form $y_p = A(x)y_1(x) + B(x)y_2(x)$

$$\begin{aligned} W(x) &= \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = \\ W_1(x) &= \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = \\ W_2(x) &= \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = \end{aligned}$$

4.
$$A'(x) = \frac{W_1}{W} =$$
$$B'(x) = \frac{W_2}{W} =$$

5. Integrating we get

$$A(x) =$$

$$B(x) =$$

6. Particular solution: $y_p =$

7. Find general solution, use constants A and B .

$$y =$$