

POUŽITÁ LITERATURA

- Achterberg, C. van, 2009: Fauna Europaea: Dacnusa sibirica. [cit. 2010-06-20]. Dostupné z WWW: <<http://www.faunaeur.org/>>.
- Albajes, R., Gullino, M. L., Lenteren, J. C., Elad, Y. (Eds.), 2000: *Integrated pest and disease management in greenhouse crops*. Springer, 568 s.
- Askew, R. R., 1968: *Hymenoptera 2. Chalcidoidea Section (b). Handbooks for the Identification of British Insects*. London, Royal entomological society. 39 s.
- Biobest, 2009: Products. [cit. 2010-06-20]. Dostupné z WWW: <<http://www.biobest.be/home/3>>.
- Biotech systems, 2008: Production: biological agents. [cit. 2010-06-20]. Dostupné z WWW: <<http://www.biotech-system.com.ua/en/production/>>.
- Canepari, C., 2004: Fauna Europaea: Cryptolaemus montrouzieri. [cit. 2010-06-20]. Dostupné z WWW: <<http://www.faunaeur.org/>>.
- Castañé, C., Alomar, O., Goula, M., Gabarra, R., 2004: Colonization of tomato greenhouses by the predatory mirid bugs Macrolophus caliginosus and Dicyphus tamaninii. *Biological Control*, 30: 591-597.
- Collyer, E., 1998: Typhlodromid mite life cycle. [cit. 2010-06-20]. Dostupné z WWW: <<http://www.hortnet.co.nz/publications/hortfacts/hf401034.htm>>.
- Driesche, R. G. van, Lyon, S. M., Hoddle, M. S., Roy,S., Sanderson, J. P., 1999: Assessment of cost and performance of Eretmocerus eremicus (Hymenoptera:aphelinidae) for whitefly (Homoptera:aleyrodidae) control in comercial poinsettia crop. *Florida Entomologist*, 82: 570-594.
- EPPO, 2008: Commercially used biological control agents - Insecta, Hymenoptera (part I). [cit. 2010-06-20]. Dostupné z WWW: <http://archives.eppo.org/EPPOStandards/biocontrol_web/hymenoptera1.htm>.
- Frank, S. D., 2009: Biological control of arthropod pests using banker plant systems: Past progress and future directions. *Biological Control*, 52: 8-16.
- Gahan, A.B., 1924: Some new parasitic Hymenoptera with notes on several described forms. *Proc. U.S. Nat. Mus.*, 4:1-23.
- Gençer, L., 2004: A study of the Chalcidoid (Hymenoptera: Chalcidoidea) Parasitoids of Leafminers (Diptera: Agromyzidae) in Ankara Province. *Turk. J. Zool.*, 28: 119-122.

- Greenberg, S. M., Jones, W.A., Liu T. 2002: Interactions Among Two Species of Eretmocerus (Hymenoptera: Aphelinidae), Two Species of Whiteflies (Homoptera: Aleyrodidae), and Tomato. *Environ. Entomol.*, 31: 397-402.
- Hayes, A. J., 1998: A laboratory study on the predatory mite, *Typhlodromus pyri* (Acarina: Phytoseiidae): II The effect of temperature and prey consumption on the numerical response of adul females. *Res. Popul. Ecol.*, 30: 13-24.
- Helyer, N., Brown, K, Cattlin, N. D., 2003: *Biological control in plant protection*. Manson publishing, London. 126 s.
- Hluchý, M., Zacharda, M., 1994: *Prostředky a systémy biologické ochrany rostlin*. Brno, Biocont Laboratory, 80 s.
- Hluchý, M., Ackermann, P., Zacharda, M., Laštůvka, Z., Bagar, M., Jetmarová, E., Vanek, G., Szöke, L, Plíšek, B., 2008: *Ochrana ovocných dřevin a révy v ekologické a integrované produkci*. Brno, Biocont Laboratory, 498 s.
- Hoddle, M. S., van Driesche, R. G., Sanderson, J. P., 1998: Biology and use of the whitefly parasitoid *Encarsia formosa*. *Annu. Rev. Entomol.*, 43: 645-669.
- Chhillar, B. S., Gulami, R., Bhatnagar, P., 2007: *Agricultural acarology*. Delhi, Daya publishing house, 355 s.
- Ibrahim, A.G., Madge, D.S., 1979: Parasitization of the chrysanthemum leaf-miner *Phytomyza syngenesiae* (Hardy) (Dipt., Agromyzidae) by *Diglyphus isaea* (Walker) (Hym. Eulophidae). *Entomologist's Monthly Magazine*, 114: 71-81.
- Karg, W., 1961: Ökologische Untersuchungen von edapischen Gamasiden (Acarina: Parasitiformes). *Pedobiologia*, 1: 53-74.
- Karg, W., 1998: Räuberisch lebende Milben als Teil des antiphytopathogenen Potentials im Boden. *Arch. Phytopath. Pflanz.*, 31: 341-347.
- Khan, I. A., Fent, M., 2004: Seasonal population dynamics of *Typhlodromus pyri* Scheuten (Acari: Phytoseiidae) in apple orchards in the region Meckenheim. *J. Pest Sci.*, 78: 1-6.
- Kobza, F. a kol. 2001. Skleníková výroba – rostlinolékařství. Mendelova zemědělská a lesnická univerzita v Brně, Brno. 65 s.
- Laan E. M, van der, Burggraaf Nierop Y. D., van, Lenteren, J.C., van, 1982: Oviposition frequency, fecundity, and life-span of *Encarsia formosa* (Hymenoptera: Aphelinidae) and *Trialeurodes vaporariorum* (Homoptera: Aleyrodidae) and migration capacity of *E. formosa* at low greenhouse temperatures. *Med. Fac. Landbouww. Rijksuniv. Gent*, 47: 511-521.

- Lenteren, J.C. van, 2008. IOBC Internet Book of Biological Control, version 5, January 2008. [cit. 2010-06-20]. Dostupné na WWW: <<http://www.unipa.it/iobc/downlaod/IOBC%20InternetBookBiCoVersion5January2008.pdf>>.
- Lenteren, J.C. van, 2003: *Quality control and production of biological control agents: theory and testing procedure*. CABI Publishing, Wallingford. 327 s.
- Lenteren, J.C. van, 1995: Integrated pest management in protected crops. In. Dent, D.R. (ed.) *Integrated Pest Management: Principles and Systems Development*, 12: 311-343. London: Chapman & Hall. 356 s.
- Lenteren, J.C. van, Szabo P, Huisman PWT., 1992: The parasite-host relationship between *Encarsia formosa* Gahan (Hymenoptera: Aphelinidae) and *Trialeurodes vaporariorum* (Westwood) (Homoptera: Aleyrodidae) XXXVII. Adult emergence and initial dispersal pattern of *E. formosa*. *J. Appl. Entomol.*, 114: 392-399.
- Lenteren, J.C. van, van Vinen A, Gast HF, Kortenhoff A., 1987: The parasite-host relationship between *Encarsia formosa* Gahan (Hymenoptera: Aphelinidae) and *Trialeurodes vaporariorum* (Westwood) (Homoptera: Aleyrodidae). XVI Food efects on oogenesis, oviposition, life-span, and fecundity of *Encarsia formosa* and other hymenopterous parasites. *J. Appl. Entomol.*, 103: 69-84.
- Lundqvist, L., 2009: Fauna Europaea: *Typhlodromus pyri*. [cit. 2010-06-20]. Dostupné z WWW: <<http://www.faunaeur.org/>>.
- McPartland, M. J., Clarke, R. C., Watson, D. P., 2000: *Hemp Diseases and Pest: Management and biological control*. CABI Publishing, Walingford. 272 s.
- Moayeri, H. R. S., Ashouri, A., Brodsgaard, H. F., Enkegaard, A., 2006: Odour-mediated preference and prey preference of *Macrolophus caliginosus* between spider mites and green peach aphids. *J. Appl. Entomol.*, 130: 504-508.
- Mohd Rasdi, Z., Fauziah, I., Wan Mohamad, W.A.K., 2009: Biology of *Macrolophus caliginosus* (Heteroptera: Miridae) Predator of *Trialeurodes vaporariorum* (Homoptera: Aleyrodidae). *International Journal of Biology*, 1: 63-70.
- Noyes, D., 2009: Fauna Europaea: *Encarsia formosa*. [cit. 2010-06-20]. Dostupné z WWW: <<http://www.faunaeur.org/>>.
- Noyes, D., 2009: Fauna Europaea: Hymenoptera: Apocrita. [cit. 2010-06-20]. Dostupné z WWW: <<http://www.faunaeur.org/>>.
- Petrova, V.; Cudare, Z., Steinite, I., 2002: The Efficiency of the Predatory Mite *Amblyseius cucumeris* (Acari: Phytoseiidae) as a Control Agent of the Strawberry Mite *Phytonemus*

- pallidus (Acari: Tarsonemidae) on Field Strawberry: Proc. 4th Int. Strawberry Symp. *Acta Hort.* 567, ISHS 2002.
- Polaszek A, Evans G.A., Bennett F.D., 1992: Encarsia parasitoids of Bemisia tabaci (Hymenoptera: Aphelinidae, Homoptera: Aleyrodidae): a preliminary guide to identification. *Bull. Entomol. Res.*, 82: 375-392.
- Qiu, Y.T., Lenteren, J.C. van, Drost, Y.C., Posthuma-Doodeman, C.J.A.M., 2004: Life-history parameters of Encarsia formosa, Eretmocerus eremicus and E. mundus, aphelinid parasitoids of Bemisia argentifolii (Hemiptera: Aleyrodidae): *Eur. J. Entomol.*, 101: 83-94.
- Rod, J., Hluchý, M., Zavadil, K., Prášil, J., Somssich, I., Zacharda, M., 2005: *Obrazový atlas chorob a škůdců zeleniny střední Evropy: Ochrana zeleniny v integrované produkci včetně prostředků biologické ochrany rostlin*. Brno, Biocont Laboratory, 392 s.
- Schelt, J. van, Mulder, S., 2000: Improved methods of testing and release of Aphidoletes aphidimyza (Diptera: Ceccidomyiidae) for aphid control in glasshouses: *Eur. J. Entomol.*, 97: 511-515.
- Sampson A. C., King V.J., 1996: Macrolophus caliginosus, field establishment and pest control effect in protected tomatoes. *IOBC/WPRS Bull.*, 19: 143-146.
- Skirvin, D.J., Fenlon, J.S., 2003: The effect of temperature on the functional response of Phytoseiulus persimilis (Acari: Phytoseiidae). *Exp. Appl. Acarol.*, 31: 37-49.
- Speyer, E. R., 1930: Biological control of the greenhouse whitefly. *Nature*, 76: 1009-1010.
- Speyer, E.R., 1929: The greenhouse whitefly (*Trialeurodes vaporariorum* Westwood). *J. R. Hort. Soc.*, 1: 181-192.
- SRS, 2010: Seznam povolených přípravků. [cit. 2010-06-20]. Dostupné z WWW: <[http://www.srs.cz/pls1/pp_public/rpg10a\\$.startup](http://www.srs.cz/pls1/pp_public/rpg10a$.startup)>.
- Sugimoto, T., Minkenberg, O.P.J.M., Takabayashi, J., Dicke, M. & Lenteren, J.C. van, 1990: Foraging for patchily-distributed leaf miners by the parasitic wasp, Dacnusa sibirica. *Res. Popul. Ecol.*, 32: 381-389.
- Šefrová, H., 2006: *Rostlinolékařská entomologie*. Brno, Konvoj, 257 s.
- Tommasini, M. G., 2003: *Evaluation of Orius species for biological control of Frankliniella occidentalis (Pergande)* (Thysanoptera: Thripidae). Thesis Wageningen university: Ponsen and Looijen b.v. Wageningen, Nizozemí. 214 s.
- Voegele, J. (ed.), 1982: *Proceedings of 1er Symposium International sur Les Trichogrammes*, 20-23 April 1982, Antibes, France. Les Colloques de l'INRA 9, Paris, 307 s.

- Weeden, C.R., Shelton, A. M., Hoffman. M. P., 1999: Biological Control: A Guide to Natural Enemies in North America. [cit. 2010-06-20]. Dostupné z WWW: <<http://www.nysaes.cornell.edu/ent/biocontrol/>>.
- Wright, E. M., Chambers, R. J., 1994: Biology of the predatory mite Hypoaspis miles (Acari: Laelapidae) , a potential biological control agent of Bradysia paupera (Dipt.: Sciaridae). *Entomophaga*, 39: 225-235.
- Zemek, R., 1993: Characteristic of development and reproduction in Typhlodromus pyri on Tetranychus urticae and Cecidophyopsis ribis. II. Progeny of overwintered females. *Exp. Appl. Acarol.*, 17: 847-858.
- Zhang, Z-Q., 2003: *Mites of Greenhouses: Identification, Biology and Control*. CABI Publishing, Wallingford, UK, 244 s.

Publikace byla realizována z grantu FRVŠ 202/2010/G4.